

UCB and Dermira enter into strategic collaboration in dermatology to broaden patient access to Cimzia[®] (certolizumab pegol)

July 3, 2014 7:00 AM CT

UCB and Dermira enter into strategic collaboration in dermatology to broaden patient access to Cimzia[®] (certolizumab pegol)

- Dermira obtains exclusive rights to develop Cimzia[®] in psoriasis in the US, Canada and the European Union
- Driven by positive Phase 2 results with Cimzia[®] in psoriasis and the Phase 3 results in psoriatic arthritis^{1,2}
- Subject to regulatory approval in psoriasis, Dermira is granted exclusive rights to market Cimzia[®] to dermatologists in the US and Canada

Brussels, Belgium and Redwood City, CA – 3rd July 2014, 0700 CEST – **regulated information** – UCB, a global biopharmaceutical leader (Euronext: UCB), and Dermira, Inc., a privately held US-based dermatology company, announced today that they have entered into an exclusive licensing agreement for the development and future commercialization of Cimzia[®] (certolizumab pegol) in dermatology. This collaboration aims to broaden patient access to Cimzia[®] and make it available to patients with psoriasis, a chronic autoimmune disorder. Cimzia[®] is not currently approved for the treatment of psoriasis by any regulatory authority.

“The Dermira team has a track record in dermatology drug development and commercialization and we look forward to our collaboration. Since the first US approval of Cimzia[®] in 2008, we continue to evaluate its potential across severe immunological diseases with the goal of bringing this treatment option to more patients,” said Professor Dr. Iris Loew-Friedrich, Chief Medical Officer and Executive Vice President, UCB. “The exploratory Phase 2 studies in patients with plaque psoriasis have shown promising data that support further clinical development.¹ In addition, the improvement in psoriasis skin symptoms observed in patients with significant skin involvement in the pivotal RAPID[®]-Psoriatic Arthritis study, were consistent with those seen in our Phase 2 study in patients with plaque psoriasis.^{1,2} The RAPID[®] study supported the approval of Cimzia[®] in psoriatic arthritis in US and EU in 2013. With Dermira, we now have a strong specialized partner with whom to move forward to make Cimzia[®] available to patients living with psoriasis and their physicians.”

“Dermira is proud to partner with UCB given their global leadership in inflammatory disease research and therapy, and product expertise with this important TNF-alpha inhibitor. If successfully developed and approved, Cimzia[®] would bring a new therapy option to adult patients with moderate to severe psoriasis. Cimzia[®] for psoriasis is our most advanced program and an important addition to our existing pipeline of dermatology products,” said Tom Wiggins, Chairman and Chief Executive Officer, Dermira.

Under the terms of the agreement, UCB grants Dermira an exclusive license to develop certolizumab pegol in psoriasis in the US, Canada and the European Union. Dermira will be responsible for Phase 3 development costs and will receive payments of up to \$49.5 million on the achievement of development and regulatory milestones. Subject to regulatory approval of Cimzia[®] in psoriasis, Dermira is granted an exclusive commercial license to market Cimzia[®] to dermatologists in the US and Canada. UCB will record the sales and Dermira will receive tiered royalty payments on those product sales which are attributable to dermatologists in the US and Canada and up to \$40 million upon the achievement of tiered commercial milestones. In support of the partnership, UCB has made a \$5 million equity investment in Dermira and a commitment to invest up to an additional \$15 million in future Dermira equity financings.

About Psoriasis

Psoriasis is a common, chronic, relapsing, immune-mediated, inflammatory disorder with primary involvement of the skin.³ It affects two to three per-cent of the world’s population – approximately 125 million people worldwide.³ Psoriasis signs and symptoms can vary from person to person but may include red patches of skin covered with silvery scales, dry, cracked skin that may bleed and thickened, pitted or ridged nails.³

In 2012, the psoriasis market (US, Japan, France, Germany, Italy, Spain and the UK) was valued at \$5.5 billion, with TNF-alpha inhibitors representing 63%.⁴

About CIMZIA[®]

CIMZIA[®] is the only Fc-free, PEGylated anti-TNF (Tumor Necrosis Factor). CIMZIA[®] has a high affinity for human TNF-alpha, selectively neutralizing the pathophysiological effects of TNF-alpha.

About CIMZIA[®] in the US⁵

In the US, CIMZIA[®] is approved for the treatment of adults with moderately to severely active rheumatoid arthritis, for the treatment of adults with active psoriatic arthritis (PsA) and for adults with active ankylosing spondylitis (AS). In addition, it is approved for reducing signs and symptoms of Crohn's disease and maintaining clinical response in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy.

Important Safety Information about CIMZIA[®] in the US

Risk of Serious Infections and Malignancy

Patients treated with CIMZIA[®] are at an increased risk for developing serious infections that may lead to hospitalization or death. Most patients who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. CIMZIA[®] should be discontinued if a patient develops a serious infection or sepsis. Reported infections include:

- Active tuberculosis, including reactivation of latent tuberculosis. Patients with tuberculosis have frequently presented with disseminated or extrapulmonary disease. Patients should be tested for latent tuberculosis before CIMZIA[®] use and during therapy. Treatment for latent infection should be initiated prior to CIMZIA[®] use.
- Invasive fungal infections, including histoplasmosis, coccidioidomycosis, candidiasis, aspergillosis, blastomycosis, and pneumocystosis. Patients with histoplasmosis or other invasive fungal infections may present with disseminated, rather than localized disease. Antigen and antibody testing for histoplasmosis may be negative in some patients with active infection. Empiric anti-fungal therapy should be considered in patients at risk for invasive fungal infections who develop severe systemic illness.
- Bacterial, viral and other infections due to opportunistic pathogens, including Legionella and Listeria.

The risks and benefits of treatment with CIMZIA[®] should be carefully considered prior to initiating therapy in patients with chronic or recurrent infection. Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with CIMZIA[®], including the possible development of tuberculosis in patients who tested negative for latent tuberculosis infection prior to initiating therapy.

Lymphoma and other malignancies, some fatal, have been reported in children and adolescent patients treated with TNF blockers, of which CIMZIA[®] is a member. CIMZIA[®] is not indicated for use in pediatric patients.

Patients treated with CIMZIA[®] are at an increased risk for developing serious infections involving various organ systems and sites that may lead to hospitalization or death. Opportunistic infections due to bacterial, mycobacterial, invasive fungal, viral, parasitic, or other opportunistic pathogens including aspergillosis, blastomycosis, candidiasis, coccidioidomycosis, histoplasmosis, legionellosis, listeriosis, pneumocystosis and tuberculosis have been reported with TNF blockers. Patients have frequently presented with disseminated rather than localized disease.

Treatment with CIMZIA[®] should not be initiated in patients with an active infection, including clinically important localized infections. CIMZIA[®] should be discontinued if a patient develops a serious infection or sepsis. Patients greater than 65 years of age, patients with co-morbid conditions, and/or patients taking concomitant immunosuppressants (e.g., corticosteroids or methotrexate) may be at a greater risk of infection. Patients who develop a new infection during treatment with CIMZIA[®] should be closely monitored, undergo a prompt and complete diagnostic workup appropriate for immunocompromised patients, and appropriate antimicrobial therapy should be initiated. Appropriate empiric antifungal therapy should also be considered while a diagnostic workup is performed for patients who develop a serious systemic illness and reside or travel in regions where mycoses are endemic.

Malignancies

During controlled and open-labeled portions of CIMZIA[®] studies of Crohn's disease and other diseases, malignancies (excluding non-melanoma skin cancer) were observed at a rate of 0.5 per 100 patient-years among 4,650 CIMZIA[®]-treated patients versus a rate of 0.6 per 100 patient-years among 1,319 placebo-treated patients. In studies of CIMZIA[®] for Crohn's disease and other investigational uses, there was one case of lymphoma among 2,657 CIMZIA[®]-treated patients

and one case of Hodgkin lymphoma among 1,319 placebo-treated patients. In CIMZIA[®] RA clinical trials (placebo-controlled and open label), a total of three cases of lymphoma were observed among 2,367 patients. This is approximately 2-fold higher than expected in the general population. Patients with RA, particularly those with highly active disease, are at a higher risk for the development of lymphoma. The potential role of TNF blocker therapy in the development of malignancies is not known.

Malignancies, some fatal, have been reported among children, adolescents, and young adults who received treatment with TNF-blocking agents (initiation of therapy ≥18 years of age), of which CIMZIA[®] is a member. Approximately half of the cases were lymphoma (including Hodgkin[™]s and non-Hodgkin[™]s lymphoma), while the other cases represented a variety of different malignancies and included rare malignancies associated with immunosuppression and malignancies not usually observed in children and adolescents. Most of the patients were receiving concomitant immunosuppressants.

Cases of acute and chronic leukemia have been reported with TNF-blocker use. Even in the absence of TNF-blocker therapy, patients with RA may be at a higher risk (approximately 2-fold) than the general population for developing leukemia.

Postmarketing cases of hepatosplenic T-cell lymphoma (HSTCL), a rare type of T-cell lymphoma that has a very aggressive disease course and is usually fatal, have been reported in patients treated with TNF blockers, including CIMZIA[®]. The majority of reported TNF blocker cases occurred in adolescent and young adult males with Crohn[™]s disease or ulcerative colitis. Almost all of these patients had received treatment with the immunosuppressants azathioprine and/or 6-mercaptopurine (6-MP) concomitantly with a TNF blocker at or prior to diagnosis. Carefully assess the risks and benefits of treatment with CIMZIA[®], especially in these patient types.

Periodic skin examinations are recommended for all patients, particularly those with risk factors for skin cancer.

Heart Failure

Cases of worsening congestive heart failure (CHF) and new onset CHF have been reported with TNF blockers. CIMZIA[®] has not been formally studied in patients with CHF. Exercise caution when using CIMZIA[®] in patients who have heart failure and monitor them carefully.

Hypersensitivity

Symptoms compatible with hypersensitivity reactions, including angioedema, dyspnea, hypotension, rash, serum sickness, and urticaria, have been reported rarely following CIMZIA[®] administration. Some of these reactions occurred after the first administration of CIMZIA[®]. If such reactions occur, discontinue further administration of CIMZIA[®] and institute appropriate therapy.

Hepatitis B Reactivation

Use of TNF blockers, including CIMZIA[®], has been associated with reactivation of hepatitis B virus (HBV) in patients who are chronic carriers of this virus. Some cases have been fatal. Test patients for HBV infection before initiating treatment with CIMZIA[®]. Exercise caution in prescribing CIMZIA[®] for patients identified as carriers of HBV, with careful evaluation and monitoring prior to and during treatment. In patients who develop HBV reactivation, discontinue CIMZIA[®] and initiate effective anti-viral therapy with appropriate supportive treatment.

Neurologic Reactions

Use of TNF blockers, including CIMZIA[®], has been associated with rare cases of new onset or exacerbation of clinical symptoms and/or radiographic evidence of central nervous system demyelinating disease, including multiple sclerosis, and with peripheral demyelinating disease, including Guillain-Barré[®] syndrome. Rare cases of neurological disorders, including seizure disorder, optic neuritis, and peripheral neuropathy have been reported in patients treated with CIMZIA[®]. Exercise caution in considering the use of CIMZIA[®] in patients with these disorders.

Hematologic Reactions

Rare reports of pancytopenia, including aplastic anemia, have been reported with TNF blockers. Medically significant cytopenia (e.g., leukopenia, pancytopenia, thrombocytopenia) has been infrequently reported with CIMZIA[®]. Advise all patients to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias or infection (e.g., persistent fever, bruising, bleeding, pallor) while on CIMZIA[®]. Consider discontinuation of CIMZIA[®] therapy in patients with confirmed significant hematologic abnormalities.

Drug Interactions

An increased risk of serious infections has been seen in clinical trials of other TNF blocking agents used in combination with anakinra or abatacept. Formal drug interaction studies have not been performed with rituximab or natalizumab; however, because of the nature of the adverse events seen with these combinations with TNF blocker therapy, similar toxicities may also result from the use of CIMZIA[®] in these combinations. Therefore, the combination of CIMZIA[®] with anakinra, abatacept, rituximab, or natalizumab is not recommended. Interference with certain coagulation assays has been detected in patients treated with CIMZIA[®]. There is no evidence that CIMZIA[®] therapy has an effect on in vivo coagulation. CIMZIA[®] may cause erroneously elevated aPTT assay results in patients without coagulation abnormalities.

Autoimmunity

Treatment with CIMZIA[®] may result in the formation of autoantibodies and, rarely, in the development of a lupus-like syndrome. Discontinue treatment if symptoms of lupus-like syndrome develop.

Immunizations

Do not administer live vaccines or live-attenuated vaccines concurrently with CIMZIA[®].

Adverse Reactions

In controlled Crohn's clinical trials, the most common adverse events that occurred in $\approx 5\%$ of CIMZIA[®] patients (n=620) and more frequently than with placebo (n=614) were upper respiratory infection (20% CIMZIA[®], 13% placebo), urinary tract infection (7% CIMZIA[®], 6% placebo), and arthralgia (6% CIMZIA[®], 4% placebo). The proportion of patients who discontinued treatment due to adverse reactions in the controlled clinical studies was 8% for CIMZIA[®] and 7% for placebo.

In controlled RA clinical trials, the most common adverse events that occurred in $\approx 3\%$ of patients taking CIMZIA[®] 200 mg every other week with concomitant methotrexate (n=640) and more frequently than with placebo with concomitant methotrexate (n=324) were upper respiratory tract infection (6% CIMZIA[®], 2% placebo), headache (5% CIMZIA[®], 4% placebo), hypertension (5% CIMZIA[®], 2% placebo), nasopharyngitis (5% CIMZIA[®], 1% placebo), back pain (4% CIMZIA[®], 1% placebo), pyrexia (3% CIMZIA[®], 2% placebo), pharyngitis (3% CIMZIA[®], 1% placebo), rash (3% CIMZIA[®], 1% placebo), acute bronchitis (3% CIMZIA[®], 1% placebo), fatigue (3% CIMZIA[®], 2% placebo). Hypertensive adverse reactions were observed more frequently in patients receiving CIMZIA[®] than in controls. These adverse reactions occurred more frequently among patients with a baseline history of hypertension and among patients receiving concomitant corticosteroids and non-steroidal anti-inflammatory drugs. Patients receiving CIMZIA[®] 400 mg as monotherapy every 4 weeks in RA controlled clinical trials had similar adverse reactions to those patients receiving CIMZIA[®] 200 mg every other week. The proportion of patients who discontinued treatment due to adverse reactions in the controlled clinical studies was 5% for CIMZIA[®] and 2.5% for placebo.

The safety profile for patients with Psoriatic Arthritis (PsA) treated with CIMZIA[®] was similar to the safety profile seen in patients with RA and previous experience with CIMZIA[®].

The safety profile for AS patients treated with CIMZIA[®] was similar to the safety profile seen in patients with RA.

For full prescribing information, please visit www.ucb.com

About CIMZIA[®] in the EU/EEA⁶

CIMZIA[®] in combination with methotrexate (MTX) is approved in the EU for the treatment of moderate to severe active RA in adult patients inadequately responsive to disease-modifying antirheumatic drugs (DMARDs) including MTX. CIMZIA[®] can be given as monotherapy in case of intolerance to MTX or when continued treatment with MTX is inappropriate. CIMZIA[®], in combination with MTX, is indicated for the treatment of active psoriatic arthritis in adults when the response to previous DMARD therapy has been inadequate. CIMZIA[®] can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate. CIMZIA[®] is also approved in the EU for the treatment of adult patients with severe active axial spondyloarthritis (axSpA) comprising:

Ankylosing spondylitis (AS) – adults with severe active AS who have had an inadequate response to, or are intolerant to non-steroidal anti-inflammatory drugs [NSAIDs].

Axial spondyloarthritis (axSpA) without radiographic evidence of AS – adults with severe active axSpA without radiographic evidence of AS but with objective signs of inflammation by elevated C-reactive protein (CRP) and/or Magnetic Resonance Imaging (MRI), who have had an inadequate response to, or are intolerant to NSAIDs.

Important Safety Information about CIMZIA® in the EU/EEA

CIMZIA® was studied in 4,049 patients with rheumatoid arthritis (RA) in controlled and open label trials for up to 92 months. The commonly reported adverse reactions (1-10%) in clinical trials with CIMZIA® and post-marketing were viral infections (includes herpes, papillomavirus, influenza), bacterial infections (including abscess), rash, headache (including migraine), asthenia, leukopenia (including lymphopenia, neutropenia), eosinophilic disorder, pain (any sites), pyrexia, sensory abnormalities, hypertension, pruritus (any sites), hepatitis (including hepatic enzyme increase), injection site reactions, and nausea. Serious adverse reactions include sepsis, opportunistic infections, tuberculosis, herpes zoster, lymphoma, leukaemia, solid organ tumours, angioneurotic oedema, cardiomyopathies (includes heart failure), ischemic coronary artery disorders, pancytopenia, hypercoagulation (including thrombophlebitis, pulmonary embolism), cerebrovascular accident, vasculitis, hepatitis/hepatopathy (includes cirrhosis), and renal impairment/nephropathy (includes nephritis). In RA controlled clinical trials, 4.4% of patients discontinued taking CIMZIA® due to adverse events vs. 2.7% for placebo.

CIMZIA® is contraindicated in patients with hypersensitivity to the active substance or any of the excipients, active tuberculosis or other severe infections such as sepsis or opportunistic infections or moderate to severe heart failure.

Serious infections including sepsis, tuberculosis and opportunistic infections have been reported in patients receiving CIMZIA®. Some of these events have been fatal. Monitor patients closely for signs and symptoms of infections including tuberculosis before, during and after treatment with CIMZIA®. Treatment with CIMZIA® must not be initiated in patients with a clinically important active infection. If an infection develops, monitor carefully and stop CIMZIA® if infection becomes serious. Before initiation of therapy with CIMZIA®, all patients must be evaluated for both active and inactive (latent) tuberculosis infection. If active tuberculosis is diagnosed prior to or during treatment, CIMZIA® therapy must not be initiated and must be discontinued. If latent tuberculosis is diagnosed, appropriate anti-tuberculosis therapy must be started before initiating treatment with CIMZIA®. Patients should be instructed to seek medical advice if signs/symptoms (e.g. persistent cough, wasting/weight loss, low grade fever, listlessness) suggestive of tuberculosis occur during or after therapy with CIMZIA®.

Reactivation of hepatitis B has occurred in patients receiving a TNF-antagonist including CIMZIA® who are chronic carriers of the virus (i.e. surface antigen positive). Some cases have had a fatal outcome. Patients should be tested for HBV infection before initiating treatment with CIMZIA®. Carriers of HBV who require treatment with CIMZIA® should be closely monitored and in the case of HBV reactivation CIMZIA® should be stopped and effective anti-viral therapy with appropriate supportive treatment should be initiated.

TNF antagonists including CIMZIA® may increase the risk of new onset or exacerbation of clinical symptoms and/or radiographic evidence of demyelinating disease; of formation of autoantibodies and uncommonly of the development of a lupus-like syndrome; of severe hypersensitivity reactions. If a patient develops any of these adverse reactions, CIMZIA® should be discontinued and appropriate therapy instituted.

With the current knowledge, a possible risk for the development of lymphomas, leukaemia or other malignancies in patients treated with a TNF antagonist cannot be excluded. Rare cases of neurological disorders, including seizure disorder, neuritis and peripheral neuropathy, have been reported in patients treated with CIMZIA®.

Adverse reactions of the hematologic system, including medically significant cytopenia, have been infrequently reported with CIMZIA®. Advise all patients to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias or infection (e.g., persistent fever, bruising, bleeding, pallor) while on CIMZIA®. Consider discontinuation of CIMZIA® therapy in patients with confirmed significant haematological abnormalities.

The use of CIMZIA® in combination with anakinra or abatacept is not recommended due to a potential increased risk of serious infections. As no data are available, CIMZIA® should not be administered concurrently with live vaccines. The

14-day half-life of CIMZIA[®] should be taken into consideration if a surgical procedure is planned. A patient who requires surgery while on CIMZIA[®] should be closely monitored for infections.

CIMZIA[®] was studied in 325 patients with active axial spondyloarthritis (axSpA) in a placebo-controlled clinical trial for up to 30 months and in 409 patients with psoriatic arthritis (PsA) in a placebo-controlled clinical trial for up to 30 months. The safety profile for axSpA and PsA patients treated with CIMZIA[®] was consistent with the safety profile in RA and previous experience with CIMZIA[®].

Please consult the full prescribing information in relation to other side effects, full safety and prescribing information. European SmPC date of revision May 2014.

http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/001037/WC500069763.pdf

References

1. Reich, K. *et al.* Successful treatment of moderate to severe plaque psoriasis with the PEGylated Fab[®] certolizumab pegol: results of a phase II, randomized, placebo controlled trial with a re-treatment extension. *British Journal of Dermatology* 2012, **167**: pp180-190
2. Mease, P. *et al.* Effect of certolizumab pegol on signs and symptoms in patients with psoriatic arthritis: 24-week results of a Phase 3 double-blind randomized placebo-controlled study (RAPID-PsA) *Ann Rheum Dis*; 2014, 73(1):48-55
3. International Federation of Psoriasis Associations. Accessed 22nd January 2014 at <http://www.worldpsoriasisday.com/web/page.aspx?refid=130>
4. Decision Resources, 2013 Immune and Inflammatory Disorders Study, Psoriasis
5. Cimzia[®] US Prescribing Information. Accessed 18th June 2014 from <http://www.ucb.com/>
6. Cimzia[®] EU Summary of Product Characteristics. Accessed 18th June 2014 from http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/001037/WC500069763.pdf

For further information, UCB:

- Antje Witte, Investor Relations UCB
T +32.2.559.9414, antje.witte@ucb.com
- Alexandra Deschner, Investor Relations, UCB
T +32 2 559 9683, alexandra.deschner@ucb.com
- Eimear O'Brien, Brand Communications, UCB
T +32.2.559.9271, eimear.obrien@ucb.com
- Andrea 2055 Woodside Road | Redwood City, CA | 94061 | Levin, US Communications, UCB
T +1.770.970.8352, andrea.levin@ucb.com
- France Nivelles, Global Communications UCB
T +32.2.559.9178, france.nivelles@ucb.com
- Laurent Schots, Media Relations, UCB
T +32.2.559.9264, laurent.schots@ucb.com

For further information, Dermira:

- Tom Wiggins, Chief Executive Officer
T: +1 650 421 7200, tom@dermira.com

About UCB

UCB, Brussels, Belgium (www.ucb.com) is a global biopharmaceutical company focused on the discovery and development of innovative medicines and solutions to transform the lives of people living with severe diseases of the immune system or of the central nervous system. With 9 000 people in approximately 40 countries, the company generated revenue of EUR 3.4 billion in 2012. UCB is listed on Euronext Brussels (symbol: UCB).

About Dermira

Dermira is a private, development-stage company focused on developing and commercializing new therapies in dermatology. Dermira was founded by an experienced management team and is advancing a pipeline of topical and systemic product candidates addressing key areas of need in dermatology. Dermira's portfolio of dermatology therapeutics includes lead product candidate Cimzia®, an Fc-free, PEGylated anti-TNF (Tumor Necrosis Factor) in clinical development for moderate to severe psoriasis; DRM04, a topical treatment for hyperhidrosis; DRM01, a topical sebum inhibitor for acne; DRM02, a topical PDE4 inhibitor for inflammatory skin diseases; and DRM05, a topical photodynamic therapy for acne. Dermira is headquartered in Redwood City, California. For more information, please visit www.dermira.com.

Forward looking statements- UCB

This press release contains forward-looking statements based on current plans, estimates and beliefs of management. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including estimates of revenues, operating margins, capital expenditures, cash, other financial information, expected legal, political, regulatory or clinical results and other such estimates and results. By their nature, such forward-looking statements are not guarantees of future performance and are subject to risks, uncertainties and assumptions which could cause actual results to differ materially from those that may be implied by such forward-looking statements contained in this press release. Important factors that could result in such differences include: changes in general economic, business and competitive conditions, the inability to obtain necessary regulatory approvals or to obtain them on acceptable terms, costs associated with research and development, changes in the prospects for products in the pipeline or under development by UCB, effects of future judicial decisions or governmental investigations, product liability claims, challenges to patent protection for products or product candidates, changes in laws or regulations, exchange rate fluctuations, changes or uncertainties in tax laws or the administration of such laws and hiring and retention of its employees. UCB is providing this information as of the date of this press release and expressly disclaims any duty to update any information contained in this press release, either to confirm the actual results or to report a change in its expectations.

There is no guarantee that new product candidates in the pipeline will progress to product approval or that new indications for existing products will be developed and approved. Products or potential products which are the subject of partnerships, joint ventures or licensing collaborations may be subject to differences between the partners. Also, UCB or others could discover safety, side effects or manufacturing problems with its products after they are marketed.

Moreover, sales may be impacted by international and domestic trends toward managed care and health care cost containment and the reimbursement policies imposed by third-party payers as well as legislation affecting biopharmaceutical pricing and reimbursement.

Forward looking statements of Dermira

This press release contains forward-looking statements based on current plans, estimates and beliefs of management. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including regarding market size and growth, product efficacy, expected legal, political, regulatory or clinical results, the achievement of development, regulatory or commercial milestones or the amount of any milestone or royalty payments, future equity financing transactions and any other projections, estimates or results. By their nature, such forward-looking statements are not guarantees of future performance and are subject to risks, uncertainties and assumptions which could cause actual results to differ materially from those that may be implied by such forward-looking statements contained in this press release. Important factors that could result in such differences include: changes in general economic, business and competitive conditions, the inability to obtain necessary regulatory approvals or to obtain them on acceptable terms, costs associated with research and development, outcomes of clinical trials, changes in the prospects for products in the pipeline or under development, the impact of competitive products and therapies, effects of future judicial decisions or governmental investigations, intellectual property and product liability claims, challenges to patent protection for products or product candidates, changes in laws or regulations, changes in financing markets, inability to raise sufficient financing, exchange rate fluctuations, changes or uncertainties in tax laws or the administration of such laws and hiring and retention of its employees. Dermira is providing this information as of the date of this press release and expressly disclaims any duty to update any information contained in this press release, either to confirm the actual results or to report a change in expectations. There is no guarantee that new product candidates in the pipeline will progress to product approval or that new indications for existing products will be developed and approved. Products or potential products which are the subject of partnerships, joint ventures or licensing collaborations may be subject to differences between the partners. Also, Dermira or others could discover safety, side effects or manufacturing problems with its products after they are marketed. Moreover, sales may be impacted by international and domestic trends toward managed care and health care cost containment and the reimbursement policies imposed by third-party payers as well as legislation affecting biopharmaceutical pricing and reimbursement.